

New Enhancement Mechanism of the Transitions in the Earth of the Solar and Atmospheric Neutrinos Crossing the Earth Core

S.T. Petcov^{a b}

^aScuola Internazionale Superiore di Studi Avanzati, and INFN (Trieste), I-34014 Trieste, Italy.

^bInst. of Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, 1784 Sofia, Bulgaria.

It is shown that the $\nu_2 \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_e$ ($\nu_e \rightarrow \nu_{\mu(\tau)}$) transitions respectively of the solar and atmospheric neutrinos in the Earth in the case of $\nu_e - \nu_{\mu(\tau)}$ mixing in vacuum, are strongly enhanced by a new type of resonance when the neutrinos cross the Earth core. The resonance is operative at small mixing angles but differs from the MSW one. It is in many respects similar to the electron paramagnetic resonance taking place in a specific configuration of two magnetic fields. The conditions for existence of the new resonance include, in particular, specific constraints on the neutrino oscillation lengths in the Earth mantle and in the Earth core, thus the resonance is a “neutrino oscillation length resonance”. It leads also to enhancement of the $\nu_2 \rightarrow \nu_e$ and $\nu_e \rightarrow \nu_s$ transitions in the case of $\nu_e - \nu_s$ mixing and of the $\bar{\nu}_\mu \rightarrow \bar{\nu}_s$ (or $\nu_\mu \rightarrow \nu_s$) transitions at small mixing angles. The presence of the neutrino oscillation length resonance in the transitions of solar and atmospheric neutrinos traversing the Earth core has important implications for current and future solar and atmospheric neutrino experiments, and more specifically, for the interpretation of the results of the Super-Kamiokande experiment.

1. Introduction

When the solar and atmospheric neutrinos traverse the Earth, the $\nu_2 \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_e$ ($\nu_e \rightarrow \nu_{\mu(\tau)}$) transitions/oscillations they undergo due to small $\nu_\mu - \nu_e$ mixing in vacuum¹ can be dramatically enhanced by a new type of resonance which differs from the MSW one and takes place when the neutrinos cross the Earth core [1]. The resonance is present in the $\nu_2 \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_e$ ($\nu_e \rightarrow \nu_{\mu(\tau)}$) transition probabilities, P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$, if the neutrino oscillation length (and mixing angles) in the Earth mantle and in the Earth core obey specific conditions [1]. When satisfied, these conditions ensure that the relevant oscillating factors in the probabilities P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ are maximal² and that this produces a resonance maximum in P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$. Accordingly, the

term “neutrino oscillation length resonance” or simply “oscillation length resonance” was used in [1] to denote the resonance of interest. There exists a beautiful analogy between the neutrino oscillation length resonance and the electron spin-flip resonance realized in a specific configuration of magnetic fields³ (see [1] for further details).

At small mixing angles ($\sin^2 2\theta \lesssim 0.05$) the maxima due to the neutrino oscillation length resonance in P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ are absolute maxima and dominate in P_{e2} and $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$: the values of the probabilities at these maxima in the simplest case of two-neutrino mixing are considerably larger - by a factor of $\sim (2.5 - 4.0)$ ($\sim (3.0 - 7.0)$), than the values of P_{e2} and $P(\nu_\mu \rightarrow \nu_e) = P(\nu_e \rightarrow \nu_{\mu(\tau)})$ at the local maxima associated with the MSW effect taking place in the Earth core (mantle). The magnitude of the enhancement due to the oscillation length resonance depends on the neutrino trajectory through the Earth core: the enhancement is maximal for the center-crossing neutrinos [1,2]. Even at small mixing angles the resonance is rel-

¹As is well-known, the $\nu_2 \rightarrow \nu_e$ transition probability accounts for the Earth effect in the solar neutrino survival probability in the case of the MSW two-neutrino $\nu_e \rightarrow \nu_{\mu(\tau)}$ and $\nu_e \rightarrow \nu_s$ transition solutions of the solar neutrino problem, ν_s being a sterile neutrino.

²Note that, in contrast, the MSW effect is a resonance amplifying the neutrino mixing.

³This analogy was brought to the attention of the author by L. Wolfenstein.

atively wide both in the neutrino energy (or resonance density) [1] - it is somewhat wider than the MSW resonance, and in the Nadir angle [2], h , specifying the neutrino trajectory in the Earth. It also exhibits strong energy dependence.

The presence of the oscillation length resonance in the transitions of solar and atmospheric neutrinos traversing the Earth has important implications [1–5] for the interpretation of the results, e.g., of the Super-Kamiokande experiment [6,7].

The Earth enhancement of the two-neutrino transitions of interest has been discussed rather extensively, see, e.g., refs. [5,8]. Some of the articles contain plots of the probabilities P_{e2} and/or $P(\nu_\mu \rightarrow \nu_e)$ or $P(\nu_e \rightarrow \nu_{\mu(\tau)})$ on which one can recognize now the dominating maximum due to the neutrino oscillation length resonance (see, e.g., [8]). However, this maximum was invariably interpreted to be due to the MSW effect in the Earth core before the appearance of [1].

2. The Neutrino Oscillation Length Resonance (NOLR)

All the interesting features of the solar and atmospheric neutrino transitions in the Earth, including those related to the neutrino oscillation length resonance, can be understood quantitatively in the framework of the two-layer model of the Earth density distribution [1,2,9,10]. The density profile of the Earth in the two-layer model is assumed to consist of two structures - the mantle and the core, having different constant densities, $\bar{\rho}_{man}$ and $\bar{\rho}_c$, and different constant electron fraction numbers, Y_e^{man} and Y_e^c ⁴. The transitions of interest of the neutrinos traversing the Earth are essentially caused by two-neutrino oscillations taking place i) first in the mantle over a distance X' with a mixing angle θ'_m and oscillation length L_{man} , ii) then in the core over a distance X'' with different mixing angle θ''_m and oscillation length L_c , and iii) again in the mantle over a distance X' with θ'_m and L_{man} . Due to the matter effect $\theta'_m, \theta''_m \neq \theta$ and $L_{man,c} \neq L_{vac}, L_{vac}$

⁴The densities $\bar{\rho}_{man,c}$ should be considered as mean densities along the neutrino trajectories. In the Earth model [11] one has: $\bar{\rho}_{man} \cong (4 - 5) \text{ g/cm}^3$ and $\bar{\rho}_c \cong (11 - 12) \text{ g/cm}^3$. For Y_e one can use the standard values [11,12] (see also [3]) $Y_e^{man} = 0.49$ and $Y_e^c = 0.467$.

being the oscillation length in vacuum (see, e.g., [13]). For fixed X' and X'' the neutrino oscillation length resonance occurs [1] if i) the relative phases acquired by the energy eigenstate neutrinos in the mantle and in the core, $\Delta E' X' = 2\pi X'/L_{man}$ and $\Delta E'' X'' = 2\pi X''/L_c$, are correlated, being odd multiples of π , so that

$$\frac{X'}{L_{man}} = k + \frac{1}{2}, \quad \frac{X''}{L_c} = k' + \frac{1}{2}, \quad (1)$$

where $k, k' = 0, 1, 2, \dots$, and if ii) the inequality

$$\cos(2\theta''_m - 4\theta'_m + \theta) (\cos(2\theta''_m - 4\theta'_m)) < 0 \quad (2)$$

is fulfilled. Condition (2) is valid for the probability P_{e2} ($P(\nu_\mu \rightarrow \nu_e)$). When equalities (1) hold, (2) ensures that P_{e2} ($P(\nu_\mu \rightarrow \nu_e)$) has a maximum. In the region of the NOLR maximum where, e.g., $\Delta E' X' \cong \pi(2k + 1)$, P_{e2} is given in the case of $\nu_e - \nu_\mu$ mixing by [1]:

$$P_{e2} \cong \sin^2 \theta + \frac{1}{4} [1 - \cos \Delta E'' X''] \times [\sin^2(2\theta''_m - 4\theta'_m + \theta) - \sin^2 \theta]. \quad (3)$$

At the NOLR maximum P_{e2} takes the form [1]

$$P_{e2}^{max} = \sin^2(2\theta''_m - 4\theta'_m + \theta). \quad (4)$$

The analogs of eqs. (3) - (4) for the probability $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ can be obtained by formally setting $\theta = 0$ while keeping $\theta'_m \neq 0$ and $\theta''_m \neq 0$ in (3) - (4). Note that one of the two NOLR requirements $\Delta E' X' = \Delta E'' X'' = \pi$ is equivalent at $\sin^2 2\theta \lesssim 0.02$ to the physical condition [1]

$$\pi \left(\frac{1}{X'} + \frac{1}{X''} \right) \cong \sqrt{2} G_F (Y_e^c \bar{\rho}_c - Y_e^{man} \bar{\rho}_{man}). \quad (5)$$

Remarkably enough, for the $\nu_2 \rightarrow \nu_e$ and $\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)}$ transitions in the Earth, the NOLR conditions (1) with $k = k' = 0$ are approximately fulfilled at small mixing angles ($\sin^2 2\theta \lesssim 0.05$) in the regions where (2) holds [1]. The associated NOLR maxima in P_{e2} and $P(\nu_\mu \rightarrow \nu_e)$ are absolute maxima (Figs. 1 - 2).

Let us note that the study performed in [1] and discussed briefly above⁵ differs substantially from

⁵For analysis of the NOLR effects in the $\nu_2 \rightarrow \nu_e$ and $\nu_e \rightarrow \nu_s$ transitions ($\nu_e - \nu_s$ mixing) and in the $\bar{\nu}_\mu \rightarrow \bar{\nu}_s$ (or $\nu_\mu \rightarrow \nu_s$) transitions at small mixing angles see [1,2].

the studies [14]. The authors of [14] considered the possibility of resonance enhancement of the $\nu_e \rightarrow \nu_{\mu(\tau)}$ transitions of neutrinos propagating in matter with density, varying *periodically* along the neutrino path (parametric resonance). It was found, in particular, that at small mixing angles strong enhancement is possible only if the neutrinos traverse at least 2 - 3 periods (in length) of the density oscillations. The density distribution in the Earth is not periodic ⁶; and in order for the oscillation length resonance ⁷ to occur periodic variation of the density is not required.

In [16] the $\nu_\mu \rightarrow \nu_s$ transitions in the Earth were considered for $\sin^2 2\theta \cong 1$. It was noticed that in the region where $\sqrt{2}G_F N_n^{man,c} \gg \Delta m^2/E$, $N_n^{man,c}$ being the neutron number density, a new maximum in $P(\nu_\mu \rightarrow \nu_s)$ appears when $\sqrt{2}G_F N_n^{man(c)} X' \cong 2\pi$, which was found to hold at $h \sim 28^\circ$. The height of the maximum is comparable to the heights of the other “ordinary” maxima present in $P(\nu_\mu \rightarrow \nu_s)$ for $\sin^2 2\theta = 1$. It is stated in [16] that the effect does not take place in the $\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)}$ transitions, which is incorrect both for $\sin^2 2\theta \ll 1$ and $\sin^2 2\theta \cong 1$ [1,2].

3. Implications of the Neutrino Oscillation Length Resonance

The implications of the oscillation length resonance enhancement of the probability P_{e2} for the Earth core crossing solar neutrinos, for the tests of the MSW $\nu_e \rightarrow \nu_{\mu(\tau)}$ and $\nu_e \rightarrow \nu_s$ solutions of the solar neutrino problem are discussed in refs. [1–4]. It is remarkable that for values of $\Delta m^2 \cong (4.0 - 8.0) \times 10^{-6} \text{ eV}^2$ from the small mixing angle (SMA) MSW solution region and the geographical latitudes at which the Super-Kamiokande, SNO and ICARUS detectors are located, the enhancement takes place in the $\nu_e \rightarrow \nu_{\mu(\tau)}$ case for values of the ^8B neutrino energy lying in the interval $\sim (5 - 12) \text{ MeV}$ to which these detectors are sensitive. The resonance max-

⁶The density change along the path of a neutrino crossing the Earth core is not periodic even in the two-layer model: it falls short of making even one and a half periods.

⁷Although this term may not be perfect, it underlines the physical essence of the new resonance. The objection to it raised in [15] is not convincing. The term “parametric resonance” used in [15], e.g., suggests incorrect analogies.




Figure 1. The probability P_{e2} as a function of h and the resonance density ρ_r for $\sin^2 2\theta = 0.01$ [2]. The prominent absolute maximum for $h \approx (0^\circ - 28^\circ)$ at $\rho_r \sim (8 - 10) \text{ g/cm}^3$ is due to the NOLR. The “shoulder” at $\rho_r \sim 13 \text{ g/cm}^3$ is caused by the MSW effect in the Earth core [1].

imum in P_{e2} at $\sin^2 2\theta = 0.01$ for the trajectory with $h = 23^\circ$, for instance, is located at $E \cong 5.3 (10.5) \text{ MeV}$ if $\Delta m^2 = 4.0 (8.0) \times 10^{-6} \text{ eV}^2$. Accordingly, at small mixing angles the NOLR is predicted [3] to produce a much bigger - by a factor of ~ 6 , day-night (D-N) asymmetry in the Super-Kamiokande sample of solar neutrino events, whose night fraction is due to the core-crossing neutrinos, in comparison with the asymmetry determined by using the *whole night* event sample. On the basis of these results it was concluded in [3] that it can be possible to test a substantial part of the MSW $\nu_e \rightarrow \nu_{\mu(\tau)}$ SMA solution region in the $\Delta m^2 - \sin^2 2\theta$ plane by performing *core* D-N asymmetry measurements. The Super-Kamiokande collaboration has already successfully applied this approach to the analysis of their solar neutrino data [6]: the limit the




Figure 2. The probability $P(\nu_{e(\mu)} \rightarrow \nu_{\mu(\tau)})$ as a function of h and $E/\Delta m^2$ for $\sin^2 2\theta = 0.01$ [2]. The absolute maximum due to the NOLR for $h \approx (0^\circ - 28^\circ)$ is clearly seen at $E/\Delta m^2 \sim (1.3 - 1.6) \times 10^6$ MeV/eV². The local maximum at $E/\Delta m^2 \sim (2.5 - 3.0) \times 10^6$ MeV/eV² is due to the MSW effect in the Earth mantle.

collaboration has obtained on the D-N asymmetry utilizing only the *core* event sample permitted to exclude a part of the MSW SMA solution region located in the area $\sin^2 2\theta \cong (0.007 - 0.01)$, $\Delta m^2 \cong (0.5 - 1.0) \times 10^{-5}$ eV², which is allowed by the mean event rate data from all solar neutrino experiments (Homestake, GALLEX, SAGE, Kamiokande and Super-Kamiokande). In contrast, the current Super-Kamiokande upper limit on the *whole night* D-N asymmetry [6] does not permit to probe the SMA solution region: the predicted asymmetry is too small [3].

The strong NOLR enhancement of the $\nu_\mu \rightarrow \nu_e$ and $\nu_e \rightarrow \nu_{\mu(\tau)}$ transitions of atmospheric neutrinos crossing the Earth core can take place at small mixing angles practically for all neutrino trajectories through the core [1], e.g., for the trajectories with $h = (0^\circ - 23^\circ)$ (Fig. 2). This

is particularly relevant for the interpretation of the results of the atmospheric neutrino experiments and for the future studies of the oscillations/transitions of atmospheric neutrinos crossing the Earth. The Super-Kamiokande collaboration has reported at this Conference strong evidences for oscillations of the atmospheric ν_μ ($\bar{\nu}_\mu$) [7]. Assuming two-neutrino mixing, the data is best described in terms of $\nu_\mu(\bar{\nu}_\mu) \leftrightarrow \nu_\tau(\bar{\nu}_\tau)$ vacuum oscillations with parameters $\Delta m^2 \cong (0.5 - 6.0) \times 10^{-3}$ eV² and $\sin^2 2\theta \cong (0.8 - 1.0)$. The possibility of two-neutrino $\nu_\mu(\bar{\nu}_\mu) \leftrightarrow \nu_e(\bar{\nu}_e)$ large mixing oscillations is disfavored by the data [7]; at $\Delta m^2 \gtrsim 2 \times 10^{-3}$ eV² it is ruled out [17].

It is a remarkable coincidence that for $\Delta m^2 \sim (0.5 - 6.0) \times 10^{-3}$ eV² and small mixing, $\sin^2 2\theta \lesssim 0.10$, the oscillation length resonance in $P(\nu_\mu \rightarrow \nu_e) = P(\nu_e \rightarrow \nu_{\mu(\tau)})$ occurs [1] for values of the energy E of the atmospheric ν_e and ν_μ which contribute either to the sub-GeV or to the multi-GeV e -like and μ -like Super-Kamiokande event samples [7]. For $\sin^2 2\theta = 0.01$, $\Delta m^2 = 5 \times 10^{-4}$; 10^{-3} ; 5×10^{-3} eV², and $h = 0^\circ$ (Earth center crossing), for instance, the absolute maximum in $P(\nu_{\mu(e)} \rightarrow \nu_{e(\mu;\tau)})$ due to the NOLR takes place at $E \cong 0.75$; 1.50; 7.5 GeV. Thus, for values of Δm^2 from the region of the $\nu_\mu \leftrightarrow \nu_\tau$ oscillation solution of the atmospheric neutrino problem, the NOLR strongly enhances the $\nu_\mu \rightarrow \nu_e$ (and $\nu_e \rightarrow \nu_{\mu(\tau)}$) transitions of the atmospheric neutrinos crossing the Earth core, making the transitions detectable even at small mixing angles. It was suggested in [1] that the excess of e -like events in the region $-1 \leq \cos \theta_z \leq -0.6$, θ_z being the Zenith angle, either in the sub-GeV or in the multi-GeV sample, observed (in both samples) in the Super-Kamiokande experiment [7], is due to $\nu_\mu \rightarrow \nu_e$ small mixing angle transitions, $\sin^2 2\theta_{e\mu} \cong (0.01 - 0.10)$, with $\Delta m^2 \sim (0.5 - 1.0) \times 10^{-3}$ eV² or respectively $\Delta m^2 \sim (2 - 6) \times 10^{-3}$ eV², strongly enhanced by the NOLR⁸. The same resonantly enhanced

⁸A more detailed investigation [2,19] performed within the indicated three-neutrino mixing scheme reveals, in particular, that the excess of e -like events in the Super-Kamiokande sub-GeV data at $-1 \leq \cos \theta_z \leq -0.6$ seems unlikely to be due to small mixing angle $\nu_\mu \rightarrow \nu_e$ transitions amplified by the oscillation length resonance.

transitions with $\Delta m^2 \sim (2 - 6) \times 10^{-3} \text{ eV}^2$ ($\Delta m^2 \sim (0.5 - 1.0) \times 10^{-3} \text{ eV}^2$) should produce at least part of the strong zenith angle dependence, exhibited by the μ -like multi-GeV (sub-GeV) Super-Kamiokande data [2].

The transitions of interest arise in a three-neutrino mixing scheme, in which the small mixing angle MSW $\nu_e \rightarrow \nu_\mu$ transitions with $\Delta m_{21}^2 \sim (4 - 8) \times 10^{-6} \text{ eV}^2$, or large mixing angle $\nu_e \leftrightarrow \nu_\mu$ oscillations with $\Delta m_{21}^2 \sim 10^{-10} \text{ eV}^2$, provide the solution of the solar neutrino problem and the atmospheric neutrino anomaly is due to $\nu_\mu \leftrightarrow \nu_\tau$ oscillations with $\Delta m_{31}^2 \sim (0.5 - 6.0) \times 10^{-3} \text{ eV}^2$ [1]. For $\Delta m_{31}^2 \gg \Delta m_{21}^2$ the three-neutrino $\nu_\mu \rightarrow \nu_e$ and $\nu_e \rightarrow \nu_{\mu(\tau)}$ transition probabilities reduce [18] to the two-neutrino transition probability $P(\nu_e \rightarrow \nu_\tau)$ (Fig. 2) with Δm_{31}^2 and $\sin^2 2\theta_{e\mu} = 4|U_{e3}|^2(1 - |U_{e3}|^2)$ playing the role of the two-neutrino oscillation parameters, where U_{e3} is the $e - \nu_3$ element of the lepton mixing matrix, ν_3 being the heaviest massive neutrino. The data [7,17] implies: $\sin^2 2\theta_{13} \lesssim 0.25$. Thus, searching for the $\nu_\mu \rightarrow \nu_e$ and $\nu_e \rightarrow \nu_{\mu(\tau)}$ transitions of atmospheric neutrinos, amplified by the oscillation length resonance, can provide also unique information about the magnitude of U_{e3} [19].

4. Conclusions

The neutrino oscillation length resonance should be present in the $\nu_2 \rightarrow \nu_e$ transitions taking place when the solar neutrinos cross the Earth core on the way to the detector, if the solar neutrino problem is due to small mixing angle MSW $\nu_e \rightarrow \nu_\mu$ transitions in the Sun. The same resonance should be operative also in the $\nu_\mu \rightarrow \nu_e$ ($\nu_e \rightarrow \nu_{\mu(\tau)}$) small mixing angle transitions of the atmospheric neutrinos crossing the Earth core if the atmospheric ν_μ and $\bar{\nu}_\mu$ indeed take part in large mixing vacuum $\nu_\mu(\bar{\nu}_\mu) \leftrightarrow \nu_\tau(\bar{\nu}_\tau)$, oscillations with $\Delta m^2 \sim (5 \times 10^{-4} - 6 \times 10^{-3}) \text{ eV}^2$, as is strongly suggested by the Super-Kamiokande data [7], and if all three flavour neutrinos are mixed in vacuum. The existence of three-flavour-neutrino mixing in vacuum is a very natural possibility in view of the present experimental evidences for oscillations/transitions of the flavour neutrinos. In both cases the oscillation length resonance produces a strong enhancement of the cor-

responding transitions probabilities, making the effects of the transitions observable even at rather small mixing angles. Actually, the resonance may have already manifested itself in the excess of e-like events at $-1 \leq \cos \theta_z \leq -0.6$ observed in the Super-Kamiokande multi-GeV atmospheric neutrino data [1,2,19]. And it can be responsible for at least part of the strong zenith angle dependence present in the Super-Kamiokande multi-GeV and sub-GeV μ -like data [2,19].

REFERENCES

1. S.T. Petcov, Phys. Lett. **B434** (1998) 321.
2. M. Chizhov, M. Maris and S.T. Petcov, Report SISSA 53/98/EP.
3. M. Maris and S.T. Petcov, Phys. Rev. **D56** (1997) 7444.
4. M. Maris and S.T. Petcov, hep-ph/9803244.
5. Q.Y. Liu, M. Maris and S.T. Petcov, Phys. Rev. **D56** (1997) 5991.
6. Y. Suzuki, these Proceedings.
7. T. Kajita, these Proceedings.
8. S.P. Mikheyev and A.Yu. Smirnov, Proc. of the Moriond Workshop on Massive Neutrinos, 1986 (eds. O. Fackler and J. Tran Thanh Van Editions Frontières, France, 1986), p. 355; A. Dar et al., Phys. Rev. **D35** (1987) 3607; M. Cribier et al., Phys. Lett. **B182** (1986) 89; A.J. Baltz and J. Weneser, Phys. Rev. **D35** (1987) 528; J.M. Gelb, W. Kwong and S.P. Rosen, Phys. Rev. Lett. **78** (1997) 2296.
9. P.I. Krastev and S.T. Petcov, Phys. Lett. **B205** (1988) 84.
10. M. Maris, Q. Liu and S.T. Petcov, study performed in December of 1996 (unpublished).
11. F.D. Stacey, *Physics of the Earth*, 2nd edition, John Wiley and Sons, New York, 1977.
12. R. Jeanloz, Annu. Rev. Earth Planet. Sci. **18** (1990) 356.
13. S.T. Petcov, hep-ph/9806466.
14. V.K. Ermilova et al., Short Notices of the Lebedev Institute **5** (1986) 26; E.Kh. Akhmedov, Yad. Fiz. **47** (1988) 475; P.I. Krastev and A.Yu. Smirnov, Phys. Lett. **B226** (1989) 341.
15. A.Yu. Smirnov, these Proceedings.
16. Q. Liu and A.Yu. Smirnov, hep-ph/9712493.
17. M. Appolonio et al. (CHOOZ Collaboration),

- Phys. Lett. **B420** (1998) 397.
18. S.T. Petcov, Phys. Lett. **B214** (1988) 259.
 19. S.T. Petcov, L. Wolfenstein and O. Yasuda,
work in progress.